




## 2025 Top 20 Protocols - After Hours Pediatric

These are the top 20 After Hours Pediatric protocols used in ClearTriage in 2025. The percentage listed after each protocol reflects the percentage of the total number of After Hours Pediatric calls that included that protocol.

- |                                      |  |
|--------------------------------------|--|
| 1. Cough – 9.4%                      | 11. Immunization Reactions – 1.8%          |
| 2. Fever – 3 Months or Older – 5.3%  | 12. Earache – 1.7%                         |
| 3. Vomiting Without Diarrhea – 4.8%  | 13. Vomiting With Diarrhea – 1.6%          |
| 4. Medication Question Call – 3.7%   | 14. Eye – Pus Or Discharge – 1.5%          |
| 5. Head Injury – 3.1%                | 15. Croup – 1.3%                           |
| 6. Sore Throat – 2.2%                | 16. Influenza (Flu) – Suspected – 1.2%     |
| 7. Rash or Redness – Widespread – 2% | 17. Rash or Redness – Localized – 1.2%     |
| 8. Diarrhea – 1.9%                   | 18. Abdominal Pain – Female – 1%           |
| 9. Colds Without Cough – 1.9%        | 19. Hand-Foot-Mouth Disease – 1%           |
| 10. Constipation – 1.8%              | 20. COVID-19 Diagnosed or Suspected – 0.9% |

### Top 20 Protocols vs All Other Calls



Percentage of total After Hours Pediatric calls using at least one of the top 20 protocols: 49.2%